

SO YOU FOUND A BABY SONGBIRD...

AWARE Wildlife Center 4158 Klondike Rd Stonecrest, GA 30038 www.awarewildlife.org

BALD W/LITTLE TO NO FEATHERS OR HAVE

BALD SPOTS NOT CAUSED BY INJURY

DO ANY OF THE FOLLOWING APPLY TO THE BIRD?

- Baby is injured, caught by cat, or caught by dog
- Baby is covered in external parasites (lice/mites/fleas)



EXAMINE THE BABY BIRD, ARE THEY...

FULLY FEATHERED, POSSIBLE BITS OF FUZZ ON THEIR HEAD, NO BALD PATCHES, AND A TAIL SHORTER THAN ADULTS











CONTACT THE AWARE HELP DESK AWARFWIIDITFF.ORG/CONTACT

or find another rehabber at awarewildlife.org/rehabbers A fledgling:

- Fully to mostly feathered w/short tail feathers 1/2" to 1" long. Beginning to learn to fly, starts from ground up.
- Still under the care of mom and dad.

If you see one:

- Look for mom and dad nearby.
 Place under shrub/cover but still in view of mom and dad.
 Keep pets out of the area/leash-walk for a few days.

EVERY EFFORT SHOULD BE MADE TO REUNITE MOM AND BABY

Try and locate nest to return baby to. It's OK to pick up the baby. The parents will NOT abandon the baby if you have touched it OR the nest.

IF YOU NEED TO BRING THE BABY INSIDE

FIRST

- Place the bird in a small, lidded box.
 - Add lining w/pillowcase/t-shirt (no strings/holes/fringe).
 - Keep in quiet, dark, warm area
- Provide a heat source. Either:
 - Microwave a sock filled with rice till warm; or
 - Heating pad, on LOW HALFWAY under the box.
 - Wrap the heat source in a towel (no holes/fringe/string) so it does not burn the baby

- During day, every 30 min:
 - Place canned dog or cat food in the baby's mouth.
 - Feed until it stops opening its mouth.
 - You should **not** have to force it, the baby should open their mouth on their own.
- - Keep baby in complete darkness and silence.
 - No feeding necessary overnight if baby kept in a dark and quiet spot.

IMPORTANT!

- If feeding the baby, please note:
 - Diet of canned dog or cat food is only nutritionally appropriate for a few days.
 Longer will cause growth abnormalities that are often fatal.
- Do not attempt to give the baby any water.
- Bring baby to a rehabber as soon as possible.

IF THE NEST CANNOT BE LOCATED/TOO HIGH UP

See the next page for more thorough information on how to create a makeshift nest!

NOTE:

IF THE BABY BECOMES WEAK, BARELY RESPONSIVE, OR STOPS POOPING FOR TWO DAYS THEN BRING INSIDE AND CONTACT A REHABBER.



SOYOU FOUND A BABY SONGBIRD...

AWARE Wildlife Center 4158 Klondike Rd Stonecrest, GA 30038 www.awarewildlife.org

MAKESHIFT NEST

TUPPERWARE NEST'



A makeshift nest that mimics most songbird nests

Round Tupperware w/holes poked in bottom (for drainage).

• Lined inside w/remains of old nest or w/dry pine straw.

Nail Tupperware Nest directly to the tree at least 6ft up.

Once nestling in position, GO INSIDE.

Mom won't return if she sees you.

• Don't be alarmed if you don't see mom.

• She only comes to nest for a few sec so that she does not draw attention from predators.

Best way to tell if baby is being care for is to:

• Check his activity levels.

Holding head up, vigorously begging for food when stimulated = doing well!

Check for poop

• Can place cardboard down to easily check in and around nest

Pooping = eating

Baby has high metabolism

One missed feeding causes drastic decrease in general status

A makeshift nest that mimic cavity nesters.

Such as woodpeckers, chickadees, and bluebirds.

Typically nest in hollowed out trees.

 Cut a door in empty jug w/holes poked in bottom (for drainage).

• Lined inside w/remains of old nest or w/dry pine straw.

Hang from a branch w/wire at least 6ft up.

Once nestling in position, GO INSIDE.

Mom won't return if she sees you.

Don't be alarmed if you don't see mom.

• She only comes to nest for a few sec so that she does not draw attention from predators.

Best way to tell if baby is being care for is to:

Chéck his activity levels.

Holding head up, vigorously begging for food when stimulated = doing well!

• Check for poop.

Can place cardboard down to easily check in and around nest.

■ Pooping = eating!

Baby has high metabolism.

 One missed feeding causes drastic decrease in general status

• If you are **unsure** if the species you found is a cavity nester, please reach out to AWARE for help with ID!

GALLON JUG NEST

